



Building Early Ontogeny Pelagic Exposure Profiles for GOA-IERP Species based on Historical Ichthyoplankton Data—Walleye Pollock




Miriam Doyle Joint Institute for the Study of the Atmosphere and Ocean, College of the Environment, University of Washington, Seattle, WA
Alaska Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA

Seasonal Patterns

Seasonal Progression in Distribution of Larvae




Vertical Distribution



- Pollock eggs occur in the plankton from late February through early May and peak in abundance during late March through early April. Larvae are most abundant from late April through early June. Larvae >15 mm are not sampled efficiently by 60 cm Bongo nets..
- Larvae up to 6 mm in length can be abundant throughout the water column to below 200 m. Larvae >6 mm are most abundant in the upper 50 m.
- Length frequency distributions indicate that growth rates of larvae are higher between May and June than between April and May.
- Larvae are abundant and ubiquitous during spring months throughout the WGOA shelf but are most abundant in Shelikof Strait. A separate source of larvae is apparent to the northeast of Kodiak Island, and in the vicinity of the Shumagins.

Interannual Patterns: Late-Spring Shelikof Time-Series



- The late-spring GOA time-series indicates periodicity in low to high levels of abundance on a scale of 2-4 years. A super-high anomaly was apparent in 1985.
- Doyle et al. (2009) have identified a positive association between late spring larval abundance and cooler winters, and strength and direction of alongshore winds during spring.
- Interannual variation in length frequency distributions of larvae reflect a slight shift in timing of sampling from year to year but also indicate diminished and enhanced growth of larvae during cold and warm years, respectively.

Interannual Variation in Larval Length Distribution

Year	Mid-Cruise Julian Day	Mid-Cruise Shift (JD)	N	WML	J-M SST	% Frequency of catch by larval length bins (mm): Walleye pollock																									WMA
						3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25			
1981	144	-4	4710	7.99																							2601.44				
1982	147	-1	407	7.43	22.56																						86.16				
1983	145.5	-2.5	2262	10.87	24.15																						351.34				
1985	148	0	2222	8.99	22.86																						46.34				
1987	142	-6	1740	7.13	25.03																						379.29				
1988	150	2	6292	8.25	24.43																						181.71				
1989	152.5	4.5	4694	8.43	19.51																						509.80				
1990	152	4	4282	9.94	20.91																						294.22				
1991	141.5	-6.5	738	6.87	19.67																						54.98				
1992	143.5	-4.5	4073	8.07	23.22																						474.51				
1993	148.5	0.5	1648	7.89	23.04																						163.88				
1994	148	0	2181	8.77	23.58																						120.31				
1995	145	-3	3535	8.18	20.99																						571.83				
1996	149	1	5050	10.29	23.21																						442.73				
1997	147	-1	3159	9.44	23.31																						548.16				
1998	146	-2	1218	8.07	28.53																						73.95				
1999	149	1	4244	7.21	20.06																						81.64				
2000	151.5	3.5	4116	8.38	23.25																						451.15				
2001	149	1	2079	8.42	25.31																						163.47				
2002	148	0	2179	6.73	23.07																						127.05				
2003	149.5	1.5	1590	8.44	27.56																						118.98				
2004	149.5	1.5	528	7.93	22.70																						18.25				
2005	148	0	1589	9.72	28.31																						64.80				
2006	147	-1	4069	8.87	23.23																						369.34				
2007	144	-4	2025	5.23	18.88																						185.19				
2008	147	-1	2168	5.94	20.15																						177.43				
2009	152.5	4.5	2772	5.96	18.69																						148.50				
2010	147	-1	3102	6.71	24.28																						423.01				
2011	156.5	8.5	1495	9.78	22.56																						179.52				

N = number of larvae measured
WMA = weighted mean abundance (no./10 m²)
WML = weighted mean length (mm)
Time-series (May 16-Jun 9) Julian Day mid-point of sampling coverage: 148
J-M SST = Cumulative mean SST for Jan through May from monthly mean values NOAA Optimum Interpolation (OI) SST V2; at Long 155.5, Lat 57.5 (color coded: light blue coldest to dark green warmest)



Correlations:
WML vs WMA: 0.08 WML vs Cum J-M SST: 0.42 WML vs Mid-Cruise Shift: 0.32
WMA vs Cum J-M SST: -0.08 WMA vs Mid-Cruise shift: -0.24

Expected tendency for smaller larvae to be more abundant than larger larvae not apparent.
Interannual variation in larval length positively related to shift in sampling dates and SST.
Interannual variation in larval abundance not related to SST, and negatively and weakly related to shift in sampling dates.

Multispecies Perspective


Early Life History (ELH) Exposure-Response Framework

Doyle and Mier 2012




- Pacific cod larvae belong to a shelf assemblage of larval fish species that prevails April-June in the WGOA (Doyle et al., 2002).
- Pacific cod, walleye pollock, and northern rock sole belong to an early life history strategy "Abundant" group that is characterized by high production of eggs/larvae with peak abundance of larvae in April, relatively limited larval duration, and moderate to high spatial ubiquity of larvae (Doyle and Mier, 2012).
- Synchronicity is observed in the abundance of these species' larvae across the time-series, with tightest synchrony between Pacific cod and northern rock sole.

Eastern vs. Western GOA



- During the historical peak period of abundance of larvae in the WGOA, walleye pollock larvae seem to be rare in the EGOA.



Synthesis

- Temporal Exposure** of walleye pollock larvae is characterized by a narrow spring peak in abundance - likely coincides with early production in GOA zooplankton.
- Spatial Exposure** extends throughout WGOA shelf waters and the major source of larvae is from spawning in Shelikof Strait, and to a lesser extent northeast of Kodiak and in the vicinity of the Shumagin Islands.
- Vertical Distribution** of larvae indicates that circulation models representing the entire water column (shelf) are suitable for tracking larval drift of newly hatched larvae, and representing the upper 50 m for larvae > 6 mm.
- The **Early Life History strategy** (ELH) common to Pacific cod, walleye pollock, and northern rock sole results in sufficient production of eggs and larvae to withstand high levels of early mortality but with limited larval duration that may risk potential mismatch with peaks in food resources, or exposure of the bulk of the larval population to other unfavorable conditions (e.g. sub-optimal temperatures and transport).
- Pelagic exposure-response coupling:** Evidence in support of this is provided by the synchrony observed among the above three species in the time-series of larval abundance and similarities in links with environmental variables (Doyle et al., 2009; Doyle and Mier, 2012.).
- Occurrence and abundance of larvae seems diminished in the EGOA relative to the WGOA.
- Gaps in ELH knowledge:**
 - Spawning areas and ELH habitat in the EGOA
 - Larval predation
 - Habitat and ecology of late stage and transitioning larvae
 - Age-0 through juvenile stage ecology (more is known than for Pacific cod)

References

Doyle, M.J., Mier, K.L., Brodeur, R.D., and Busby, M.S. (2002). Regional variations in springtime ichthyoplankton assemblages in the northeast Pacific Ocean. *Prog. Oceanogr.* 53(2-4): 247-282.

Doyle, M.J., Picquelle, S.J., Mier, K.L., Spillane, M.C., and Bond, N.A. (2009). Larval fish abundance and physical forcing in the Gulf of Alaska, 1981-2003. *Prog. Oceanogr.* 80:163-187.

Doyle, M.J. and Mier, K.L. (2012). A new conceptual framework for evaluating the early ontogeny phase of recruitment processes among marine fish species. *Can. J. Fish. Aquat. Sci.* 69: 2112-2129.